

Thromboelastometry and Thrombelastography Analysis under Normal Physiological Conditions – Systematic Review

Marcel Adler^a Sandra Ivic^b Nicolas S. Bodmer^b Hugo ten Cate^c Lucas M. Bachmann^b
Walter A. Wuillemin^d Michael Nagler^{a,d}

^a Department of Hematology and Central Hematology Laboratory, Inselspital, Bern University Hospital, and Department of Clinical Research, University of Bern, Bern, Switzerland;

^b medignition Inc, Research Consultants, Zurich, Switzerland;

^c Laboratory of Clinical Thrombosis and Hemostasis, and Cardiovascular Research Institute, Maastricht University Medical Center, Maastricht, The Netherlands;

^d Division of Hematology and Central Hematology Laboratory, Luzerner Kantonsspital, University of Bern, Lucerne, Switzerland

Keywords

Thromboelastometry · Thrombelastography · Technical efficacy · Healthy subjects · Hemostasis

Summary

Background: Studies investigating thromboelastometry or thrombelastography analyses in a physiological context are scattered and not easy to access. **Objective:** To systematically retrieve and describe published reports studying healthy subjects and targeting at the correlation of ROTEM[®] and TEG[®] measurements with conventional parameters of hemostasis. **Methods:** Systematic Review: Papers were searched in Medline, Scopus and the Science Citation Index database. Reference lists of included studies and of reviews were screened. To be included papers had to report ROTEM or TEG data on healthy subjects. Two reviewers screened papers for inclusion, read full texts of potentially relevant papers, and extracted data of included papers. **Results:** Searches identified 1,721 records of which 1,713 were either excluded immediately or after reading the full text. The remaining 8 studies enrolled 632 subjects. The association of conventional parameters of hemostasis with ROTEM and with TEG was investigated in one and two studies, respectively. Overall correlation was limited and ranged from 0.0 to 0.40 (total thrombus generation vs. fibrino-

gen; clotting time INTEM vs. activated partial thromboplastin time). **Conclusions:** Studies assessing the relationship between thromboelastometry or thrombelastography analyses and conventional parameters of hemostasis in healthy subjects remains scarce, and correlations are limited. Further research is needed to understand the physiology of thromboelastometry and thromboelastography parameters.

© 2017 S. Karger GmbH, Freiburg

Introduction

Back in 1978, Loop and Lusted [1] proposed the first phased evaluation of new medical tests. To date, after the publication of at least 18 additional recommendations, (reviewed in [2]) there is a broad consensus that first evaluations should address parameters of technical efficacy such as minimal detection level, circadian fluctuation, reproducibility, and correlation with established tests in healthy volunteers in order to understand the physiology of the analyzed parameters. It has been recognized that sophisticated and expensive tests that are disseminated without suitable evaluations can subsequently be found to have marginal clinical value and economic benefit. Well known examples include the carcinoembryonic antigen test in the diagnosis of colon cancer [3], iodine-125-labeled fibrinogen scans in the diagnosis of deep venous

Marcel Adler and Sandra Ivic contributed equally.

thrombi [4], or rapid magnetic resonance imaging in the management of patients with low back pain [5].

Thromboelastometry (ROTEM®) and thrombelastography (TEG®) analysis, two methods evaluating hemostasis, are increasingly used in clinical practice, based on the anticipation that they improve the management of acute bleeding [6, 7]. Several possible advantages over conventional laboratory parameters promote their use in emergency units and operating rooms. The analyzer is easy to use, and tests are readily available. Results are displayed graphically, allowing an intuitive interpretation [8]. Furthermore, tests claim to provide global information on all aspects of hemostasis, including fibrinogen level, platelet function, coagulation cascade, cross-linking of fibrin, and fibrinolysis. In addition, whereas the therapeutic consequences of results obtained with conventional parameters often remains uncertain, thromboelastometry/thromboelastography results suggest specific interventions to enhance hemostasis immediately [9]. Several studies explored their role in the detection of coagulopathies and changes in bleeding management as well as in the perioperative setting [10–14]. However, despite a vast amount of investigations, the clinical value of these methods has been challenged repeatedly [15–19]. It has been argued that the lack of rigorous examinations assessing the correlation of thromboelastometry and thromboelastography measurements with conventional parameters of hemostasis jeopardizes its clinical interpretation and usefulness [8, 20, 21].

To the best of our knowledge, however, there is no systematic review assessing the available evidence on thromboelastometry and thromboelastography examinations under physiological conditions in healthy volunteers. We therefore set out to systematically search, describe and inventory published reports on ROTEM and TEG data in healthy subjects, targeting at the correlation of thromboelastometry and thromboelastography measurements with conventional parameters of hemostasis.

Material and Methods

The present systematic reviews was done according to the PRISMA statement [22].

Search Strategy

To identify papers investigating correlations of ROTEM and TEG measurements with conventional parameters of hemostasis under physiological conditions, we searched (PRE-)MEDLINE (PubMed interface) using a search string including the two Medical Subject Headings (MeSH) ‘Thrombelastography/methods’[Mesh], ‘Blood Coagulation Tests’[Mesh] and the free text terms thromboelastography, thromboelastometry, ROTEM, and TEG. SCOPUS was searched using the following algorithm: TITLE-ABS-KEY(thromboelastography OR thromboelastometry) AND (valid*) AND (LIMIT-TO(DOCTYPE, ‘ar’)) AND (LIMIT-TO(SUBJAREA, ‘MEDI’)). We also searched the Web of Science database entering 4 seminal papers [8, 23–25] and checked their citations for potentially relevant papers. Searches were complemented checking reference lists of included studies and of a recently published Cochrane review [15]. Electronic searches were done from inception to November 2016.

Selection Criteria for Inclusion

To be included a paper had to use whole blood or citrated blood obtained from healthy volunteers and had to assess either ROTEM or TEG. Additional

criteria had to be fulfilled for inclusion of correlation studies: ROTEM and TEG data in combination with at least one established parameter of hemostasis such as activated partial thromboplastin time (aPTT), prothrombin time (PT), platelet count (PLT), fibrinogen level, or d-dimers.

Exclusion Criteria

We excluded papers assessing trauma patients, patients undergoing cardiac surgery, and other patients with severe illness as well as studies including pregnant women, children, or patients with sepsis if the study did not include any kind of control arm with healthy subjects.

Head-to-head comparisons between ROTEM and TEG were also excluded, if they were assessed in absence of an established hemostasis method.

Selection Process and Data Abstraction

In the case of multiple publications on the same participants, the most complete report was chosen for each study. We extracted data in duplicate, and a third reviewer resolved any discrepancies if the two reviewers disagreed. Of each study we extracted participants’ age, number of female participants, the total number of included subjects, the device employed (ROTEM, TEG), and the hemostasis parameter assessed. In case of TEG studies these were reaction time (R time), time to maximum clot formation (K time), α angle, MA (maximal amplitude), SEMS (shear elastic modulus strength), CI (coagulation index), MTG (maximal thrombus generation), TMG (time to maximal thrombus generation), and TTG (total thrombus generation). In the ROTEM studies CT (clotting time), CFT (clot formation time), α angle, and MCF (maximum clot firmness) for both INTEM and EXTEM assay were recorded. Finally, we also extracted which of the established methods was used as the comparator test and what type of correlation or concordance parameters was used.

Thromboelastometry/Thromboelastography Analysis

The analytical principle of thromboelastometry and thromboelastography analysis is discussed in detail elsewhere [13, 20, 26–28]. In brief, the viscoelastic properties of a forming clot are assessed by recording the oscillations of a pin or a cup containing a citrated whole blood sample. Different aspects of clot formation are represented as parameters of the analysis. For example, the time from addition of the reagent until start of clot formation is denoted as CT in case of ROTEM, and R time in case of TEG; the clot strength is reported as MCF, or MA. Parameters are measured after addition of different activators. By analogy with PT, tissue factor reagent is added in case of EXTEM (ROTEM) and Rapid-TEG. Comparable to the aPTT, contact phase activators are added in case of INTEM (ROTEM) or standard TEG.

Results

Selection Process


Searches retrieved 1,721 records of which 1,588 had to be excluded after screening title or abstract. Full texts of 133 articles were examined for inclusion. Of these, 125 articles fulfilled at least one exclusion criterion leaving 8 articles for detailed examination [24, 29–35]. The selection process is described in figure 1.

Included Studies

Thromboelastometry or thromboelastography measurements in healthy volunteers have been performed in 8 studies, enrolling 632 subjects (table 1) [24, 29–35]. Five studies reported thromboelastometry/thromboelastography results but did not investigate correlations with conventional parameters of hemostasis [24, 29, 30, 34, 35]. Lang and colleagues [24] conducted a multicenter study to establish reference values for ROTEM thromboelastome-

Table 1. Characteristics of identified evaluation studies in healthy volunteers

Author, year	Mean age (SD), years	Age range, years	Female, n (%)	Total (n)	Device	Correlation/concordance
Lang et al., 2005 [24]	45.4 (17.6); 43.1 (15.9); 38.8 (14.4)	–	161 (61.5%)	262	ROTEM	–
Rivard et al., 2005 [32]	–	–	2 (50%)	4	TEG	linear/non-linear regression
Tripodi et al., 2009 [35]	–	20–60	28 (48%)	58	ROTEM	–
Huissoud et al., 2009 [30]	30 (median)	26–33	20 (100%)	20	ROTEM	–
Scarpelini et al., 2009 [34]	36.8 (11.2)	–	40 (34%)	118	TEG	–
Roeloffzen et al., 2010 [33]	49.5 (25.5)	19–87	60 (50%)	120	TEG	Pearson's correlation coefficients
Foley et al., 2012 [29]	–	29–32	0 (0%)	5	TEG	–
Kim et al., 2013 [31]	44	22–71	19 (42%)	45	ROTEM	Pearson's correlation / linear regression

**Fig. 1.** Study flow.

try. 262 individuals were included in five centers in Germany, Austria, and France. Even though considerable variation between centers existed, 'orientating reference ranges' were calculated for INTEM, EXTEM, and FIBTEM parameters (CT, CFT, α angle, A10, A20, A30, MCF, CLI30, and ML). Tripodi and co-workers [35] conducted ROTEM measurements in 58 healthy volunteers to establish reference ranges for an observation study in cirrhosis patients. However, results were not reported, and no correlation studies have been done. 20 healthy women were studied using ROTEM in an investigation by Huissoud and colleagues [30], serving as a control for thromboelastometry patterns in pregnancy. Median values and inter-quartile ranges of EXTEM, INTEM, FIBTEM, and APTEM parameters were reported. TEG was conducted in 5 volunteers in a study by Foley et al. [29] aiming to improve the thromboelastography assay, and individual measurements were reported.

Correlation with Conventional Parameters of Hemostasis

Three studies reported the extent of association between thromboelastometry/thromboelastography parameters and conventional coagulation assays [31–33]. The details are reported in table 2. Two studies focused on thromboelastography [32, 33]. In a study enrolling 120 participants (50% female) of 49.5 years on average, Roeloffzen and colleagues [33] compared R time with the aPTT, K time with the hemoglobin level, aPTT and the fibrinogen level, α angle with aPTT and the fibrinogen level, MA with the fibrinogen level, SEMS with the fibrinogen level, the CI with aPTT and the fibrinogen level, MTG with the fibrinogen level, TMG with aPTT, and the TTG with the fibrinogen level. Pearson's correlation coefficient was reported and varied between 0.21 (TMG vs. aPTT) and 0.40 (TTG vs. fibrinogen level). Rivard and co-workers [32] studied the association between a TEG parameter used as surrogate for TTG with thrombin generation as measured using thrombin/antithrombin complexes (TAT) in 4 volunteers. A correlation coefficient of 0.85 was determined from a linear regression and a correlation coefficient of 0.94 from a linear regression using the ln of TAT.

Associations between thromboelastometry parameters and conventional parameters of hemostasis were investigated in one study only [31]. Kim and co-workers [31] correlated CT, CFT, α angle, CFR, and MCF with PT as well as aPTT. No significant correlation was found between any EXTEM or INTEM parameters and PT. For INTEM, the CT value was significantly correlated with aPTT only ($r = 0.41$).

Discussion

Main Findings

Using comprehensive retrieval methods, our systematic review only found few studies assessing thromboelastometry as well as thromboelastography and correlations thereof with established parameters of hemostasis in healthy volunteers. The studies revealed only a weak correlation.

Findings in Context

There are several studies investigating the relationship of thromboelastometry or thromboelastography in animals, e.g.,

Table 2. Correlation of thromboelastometry/thrombelastography parameters with conventional parameters of hemostasis in healthy volunteers

Author, year	n	Device	Test	Pearson's correlation coefficient			
				Hb	aPTT	PT	fibrinogen
Roeloffzen et al., 2010 [33]	120	TEG	R time	NS	0.36 (p < 0.05)		NS
	120	TEG	K time	0.24 (p < 0.05)	0.24 (p < 0.05)		–0.27 (p < 0.05)
	120	TEG	α angle	NS	–0.22 (p < 0.05)		–0.28 (p < 0.05)
	120	TEG	MA	NS	NS		0.37 (p < 0.05)
	120	TEG	SEMS	NS	NS		0.37 (p < 0.05)
	120	TEG	CI	NS	–0.21 (p < 0.05)		0.25 (p < 0.05)
	120	TEG	MTG	NS	NS		0.27 (p < 0.05)
	120	TEG	TMG	NS	0.21 (p < 0.05)		NS
	120	TEG	TTG	NS	NS		0.40 (p < 0.05)
	120	TEG					
Rivard et al., 2005 [32]	4	TEG	TTG	–	–		0.85 [#] / 0.94 [†] (p < 0.05)
Kim et al., 2013 [31]	45	ROTEM	CT EXTEM	–	–0.223 (p = 0.17)	–0.137 (p = 0.46)	–
	45	ROTEM	CFT EXTEM	–	–0.149 (p = 0.36)	–0.189 (p = 0.24)	–
	45	ROTEM	α angle EXTEM	–	0.079 (p = 0.63)	0.005 (p = 0.89)	–
	45	ROTEM	MCF EXTEM	–	0.005 (p = 0.97)	–0.164 (p = 0.31)	–
	45	ROTEM	CT INTEM	–	0.406 (p = 0.009)	0.12 (p = 0.46)	–
	45	ROTEM	CFT INTEM	–	–0.12 (p = 0.46)	–0.194 (p = 0.23)	–
	45	ROTEM	α angle INTEM	–	–0.053 (p = 0.75)	0.122 (p = 0.45)	–
	45	ROTEM	MCF INTEM	–	–0.014 (p = 0.93)	0.003 (p = 0.98)	–
	45	ROTEM					
	45	ROTEM					

^aPTT = Activated partial thromboplastin time; CI = coagulation index; CT = clotting time; CFT = clot formation time; Hb = hemoglobin level; K time = time to maximum clot formation; MA = maximal amplitude; MCF = maximum clot firmness; MTG = maximal thrombus generation; NS = not stated; PT = prothrombin time; R time = reaction time; SEMS = shear elastic modulus strength; TAT thrombin/antithrombin complex; TMG = time to maximal thrombus generation; TTG = total thrombus generation.

[#]Derived from linear regression analyses.

[†]Derived from log-linear regression analysis (ln(TAT)).

Mauch et al. [36]. In this study the intrarater and interrater validity of the ROTEM data was analyzed with blood from healthy pigs. Another prospective study compared the coagulation profiles of healthy foals using standard techniques (PT, aPTT, fibrinogen concentration, and antithrombin) with the TEG analyzer [37]. The validity of the ROTEM and TEG was verified through comparison with standard methods in the context of specific circumstances, as for example orthotopic liver transplantation [38–40], during aortic surgery [12, 41, 42], or in trauma patients [18, 43, 44]. We also found a few trials reporting on head-to-head comparisons between ROTEM and TEG [21, 45, 46]. Although these approaches also yield valuable information, we think that they do not make comparisons with standardized methods in healthy subjects superfluous.

Strength and Limitations

We applied up-to-date and rigorous systematic review methods to retrieve and assess the available evidence. In view of the risk that potentially relevant studies could be missed due to ambiguous indexing in the various databases, we applied an over-inclusive approach. These sensitive searches retrieved over 1,700 records. Screening for inclusion was made in duplicate to reduce the risk to miss relevant papers. We expected to find a reasonable number of studies that would allow us to provide quantitative summaries of various comparisons. Moreover, out of the many available, we hoped to depict those parameters from thromboelastometry and thromboelastography that would be most suited for investigations in various clinical settings such as acute bleeding, applications in emergency rooms, and for treatment of patients with specific hemostatic deficiencies. Despite our motivation, the current evidence base impeded us from generating meaningful summaries for further research and clinical practice.

Implications for Clinical Practice

The small number of studies examining thromboelastometry or thromboelastography in a cohort of healthy volunteers and systematically comparing the so obtained findings to conventional assays of hemostasis is remarkable. One may argue that the clinical

value of established coagulation parameters is limited as well. It is true that results of coagulation parameters such as PT not directly prompt any specific treatment. Nevertheless, they are still important biomarkers for the detection of coagulopathies in the setting of acute bleeding [47–49].

Implications for Research

Following existing guidelines of test evaluation (reviewed and discussed in [2]), we propose that further research should first investigate the distribution of normal values of ROTEM and TEG parameters in reasonably sized studies enrolling healthy subjects. Second, distributions should be assessed in groups of patients with a specific illness and compared to distributions of healthy subjects. In a third step, assessing test performance characteristics such as sensitivity and specificity should be performed, taking contextual clinical information into account. Fourth, studies should determine the added value of thromboelastometry/thromboelastography in context of other clinical information about the patients' state that is available prior testing. Finally, decision-analytic models should be performed determining the optimal areas for clinical use, taking clinical outcomes into account.

Conclusions

Studies assessing the relationship between thromboelastometry or thromboelastography analyses and parameters of hemostasis in healthy subjects remains scarce. From a hemostaseologic standpoint, further physiologic research is needed to elucidate the significance of thromboelastometry and thromboelastography for clinical practice and research.

Disclosure Statement

Financial support for the submitted work was received from the Research Fund Haematology Cantonal Hospital Lucerne. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. MN has received research grants, lecture fees, or consultancy fees from Bayer, Roche Diagnostics, Instrumentation Laboratory, Boehringer Ingelheim, and CSL Behring.

References

- 1 Loop JW, Lusted LE: American College of Radiology Diagnostic Efficacy Studies. *AJR Am J Roentgenol* 1978;131:173–179.
- 2 Lijmer JG, Leeflang M, Bossuyt PM: Proposals for a phased evaluation of medical tests. *Med Decis Making* 2009;29:E13–21.
- 3 Fletcher RH: Carcinoembryonic antigen. *Ann Intern Med* 1986;104:66–73.
- 4 Lensing AW, Hirsh J: 125I-fibrinogen leg scanning: reassessment of its role for the diagnosis of venous thrombosis in post-operative patients. *Thromb Haemost* 1993;69:2–7.
- 5 Hollingworth W, Gray DT, Martin BI, Sullivan SD, Deyo RA, Jarvik JG: Rapid magnetic resonance imaging for diagnosing cancer-related low back pain. *J Gen Intern Med* 2003;18:303–312.
- 6 Fassl J, Matt P, Eckstein F, Filipovic M, Gregor M, Zenklusen U, Seeberger MD, Bolliger D: Transfusion of allogeneic blood products in proximal aortic surgery with hypothermic circulatory arrest: effect of thromboelastometry-guided transfusion management. *J Cardiothorac Vasc Anesth* 2013;27:1181–1188.
- 7 Kozek-Langenecker SA, Afshari A, Albaladejo P, Santullano CA, De Robertis E, Filipescu DC, Fries D, Grolinger K, Haas T, Imberger G, Jacob M, Lance M, Llau J, Mallett S, Meier J, Rahe-Meyer N, Samama CM, Smith A, Solomon C, Van der Linden P, Wikkelsso AJ, Wouters P, Wyffels P: Management of severe perioperative bleeding: guidelines from the European Society of Anaesthesiology. *Eur J Anaesthesiol* 2013;30:270–382.
- 8 Luddington RJ: Thromboelastography/thromboelastometry. *Clin Lab Haematol* 2005;27:81–90.
- 9 Schochl H, Nienaber U, Hofer G, Voelckel W, Jambor C, Scharbert G, Kozek-Langenecker S, Solomon C: Goal-directed coagulation management of major trauma patients using thromboelastometry (ROTEM)-guided administration of fibrinogen concentrate and prothrombin complex concentrate. *Crit Care* 2010;14:R55.
- 10 Fenger-Eriksen C, Ingerslev J, Tonnesen E, Sorensen B: Citrate artificially masks the haemostatic effect of recombinant factor VIIa in dilutional coagulopathy. *Ann Hematol* 2009;88:255–260.
- 11 Fenger-Eriksen C, Jensen TM, Kristensen BS, Jensen KM, Tonnesen E, Ingerslev J, Sorensen B: Fibrinogen substitution improves whole blood clot firmness after dilution with hydroxyethyl starch in bleeding patients undergoing radical cystectomy: a randomized, placebo-controlled clinical trial. *J Thromb Haemost* 2009;7:795–802.

- 12 Girdauskas E, Kempfert J, Kuntze T, Borger MA, Enders J, Fassl J, Falk V, Mohr FW: Thromboelastometrically guided transfusion protocol during aortic surgery with circulatory arrest: a prospective, randomized trial. *J Thorac Cardiovasc Surg* 2010;140:1117–1124.
- 13 Rugeri L, Levrat A, David JS, Delecroix E, Floccard B, Gros A, Allaouchiche B, Negrier C: Diagnosis of early coagulation abnormalities in trauma patients by rotation thrombelastography. *J Thromb Haemost* 2007;5: 289–295.
- 14 Shore-Lesserson L, Manspeizer HE, DePerio M, Francis S, Vela-Cantos F, Ergin MA: Thromboelastography-guided transfusion algorithm reduces transfusions in complex cardiac surgery. *Anesth Analg* 1999; 88:312–319.
- 15 Afshari A, Wikkelsø A, Brok J, Møller AM, Wetterslev J: Thrombelastography (TEG) or thromboelastometry (ROTEM) to monitor haemotherapy versus usual care in patients with massive transfusion. *Cochrane Database Syst Rev* 2011;3:CD007871.
- 16 Coakley M, Hall JE, Evans C, Duff E, Billing V, Yang L, McPherson D, Stephens E, Macartney N, Wilkes AR, Collins PW: Assessment of thrombin generation measured before and after cardiopulmonary bypass surgery and its association with postoperative bleeding. *J Thromb Haemost* 2011;9:282–292.
- 17 Lee GC, Kicza AM, Liu KY, Nyman CB, Kaufman RM, Body SC: Does rotational thromboelastometry (ROTEM) improve prediction of bleeding after cardiac surgery? *Anesth Analg* 2012;115:499–506.
- 18 Raza I, Davenport R, Rourke C, Platten S, Manson J, Spoor C, Khan S, De'Ath HD, Allard S, Hart DP, Pasi KJ, Hunt BJ, Stanworth S, MacCallum PK, Brohi K: The incidence and magnitude of fibrinolytic activation in trauma patients. *J Thromb Haemost* 2013;11:307–314.
- 19 Wikkelsøe AJ, Afshari A, Wetterslev J, Brok J, Moeller AM: Monitoring patients at risk of massive transfusion with thrombelastography or thromboelastometry: a systematic review. *Acta Anaesthesiol Scand* 2011;55: 1174–1189.
- 20 Chitlur M, Lusher J: Standardization of thromboelastography: values and challenges. *Semin Thromb Hemost* 2010;36:707–711.
- 21 Chitlur M, Sorensen B, Rivard GE, Young G, Ingerslev J, Othman M, Nugent D, Kenet G, Escobar M, Lusher J: Standardization of thromboelastography: a report from the TEG-ROTEM working group. *Haemophilia* 2011;17:532–537.
- 22 Moher D, Liberati A, Tetzlaff J, Altman DG: Preferred reporting items for systematic reviews and meta-analyses: the PRISMA Statement. *Open Med* 2009;3:e123–130.
- 23 Di Benedetto P, Baciarello M, Cabetti L, Martucci M, Chiaschi A, Bertini L: Thrombelastography. Present and future perspectives in clinical practice. *Minerva Anestesiologica* 2003;69:501–509, 509–515.
- 24 Lang T, Bauters A, Braun SL, Potzsch B, von Pape KW, Kolde HJ, Lakner M: Multi-centre investigation on reference ranges for ROTEM thromboelastometry. *Blood Coagul Fibrinolysis* 2005;16:301–310.
- 25 Salooja N, Perry DJ: Thrombelastography. *Blood Coagul Fibrinolysis* 2001;12:327–337.
- 26 Lang T, von Depka M: Possibilities and limitations of thrombelastometry/-graphy (in German). *Hamostaseologie* 2006;26(suppl):S20–29.
- 27 Nagler M, ten Cate H, Kathriner S, Casutt M, Bachmann LM, Willemin WA: Consistency of thromboelastometry analysis under scrutiny: results of a systematic evaluation within and between analysers. *Thromb Haemost* 2014;111:1161–1166.
- 28 Nagler M, Kathriner S, Bachmann LM, Willemin WA: Impact of changes in haematocrit level and platelet count on thromboelastometry parameters. *Thromb Res* 2013;131:249–253.
- 29 Foley JH, Butenas S, Mann KG, Brummel-Ziedins KE: Measuring the mechanical properties of blood clots formed via the tissue factor pathway of coagulation. *Anal Biochem* 2012;422:46–51.
- 30 Huisoud C, Carrabin N, Benchaib M, Fontaine O, Levrat A, Massignon D, Touzet S, Rudigoz RC, Berland M: Coagulation assessment by rotation thromboelastometry in normal pregnancy. *Thromb Haemost* 2009;101:755–761.
- 31 Kim B, Quan ML, Goh R, Kim JE, Woo KS, Kim MH, Han JY: Comparison of prolonged prothrombin and activated partial thromboplastin time results with thrombelastograph parameters. *Lab Med* 2013;44:319–323.
- 32 Rivard GE, Brummel-Ziedins KE, Mann KG, Fan L, Hofer A, Cohen E: Evaluation of the profile of thrombin generation during the process of whole blood clotting as assessed by thrombelastography. *J Thromb Haemost* 2005;3:2039–2043.
- 33 Roeloffzen WW, Kluin-Nelemans HC, Mulder AB, Veeger NJ, Bosman L, de Wolf JT. In normal controls, both age and gender affect coagulability as measured by thrombelastography. *Anesth Analg* 2010;110:987–994.
- 34 Scarpellini S, Rhind SG, Nascimento B, Tien H, Shek PN, Peng HT, Huang H, Pinto R, Speers V, Reis M, Rizoli SB: Normal range values for thrombelastography in healthy adult volunteers. *Braz J Med Biol Res* 2009; 42:1210–1217.
- 35 Tripodi A, Primignani M, Chantarangkul V, Viscardi Y, Dell'Era A, Fabris FM, Mannucci PM: The coagulopathy of cirrhosis assessed by thromboelastometry and its correlation with conventional coagulation parameters. *Thromb Res* 2009;124:132–136.
- 36 Mauch J, Spielmann N, Hartnack S, Madjdpour C, Kutter AP, Bettschart-Wolfensberger R, Weiss M, Haas T: Intrarater and interrater variability of point of care coagulation testing using the ROTEM delta. *Blood Coagul Fibrinolysis* 2011;22:662–666.
- 37 Mendez-Angulo JL, Mudge M, Zaldivar-Lopez S, Villar-Saavedra P, Couto G: Thromboelastography in healthy, sick non-septic and septic neonatal foals. *Aust Vet J* 2011;89:500–505.
- 38 Coakley M, Reddy K, Mackie I, Mallett S: Transfusion triggers in orthotopic liver transplantation: a comparison of the thromboelastometry analyzer, the thrombelastogram, and conventional coagulation tests. *J Cardiothorac Vasc Anesth* 2006;20:548–553.
- 39 Herbstreit F, Winter EM, Peters J, Hartmann M: Monitoring of haemostasis in liver transplantation: comparison of laboratory based and point of care tests. *Anaesthesia* 2010;65:44–49.
- 40 Rouillet S, Pillot J, Freyburger G, Biais M, Quinart A, Rault A, Revel P, Sztark F: Rotation thromboelastometry detects thrombocytopenia and hypofibrinogenemia during orthotopic liver transplantation. *Br J Anaesth* 2010;104:422–428.
- 41 Ogawa S, Szlam F, Chen EP, Nishimura T, Kim H, Roback JD, Levy JH, Tanaka KA: A comparative evaluation of rotation thromboelastometry and standard coagulation tests in hemodilution-induced coagulation changes after cardiac surgery. *Transfusion* 2011;52: 14–22.
- 42 Solomon C, Cadamuro J, Ziegler B, Schochl H, Varvenne M, Sorensen B, Hochleitner G, Rahe-Meyer N: A comparison of fibrinogen measurement methods with fibrin clot elasticity assessed by thromboelastometry, before and after administration of fibrinogen concentrate in cardiac surgery patients. *Transfusion* 2011; 51:1695–1706.
- 43 Davenport R, Manson J, De'Ath H, Platten S, Coates A, Allard S, Hart D, Pearce R, Pasi KJ, MacCallum P, Stanworth S, Brohi K: Functional definition and characterization of acute traumatic coagulopathy. *Crit Care Med* 2011;39:2652–2658.
- 44 Tauber H, Innerhofer P, Breitkopf R, Westermann I, Beer R, El Attal R, Strasak A, Mittermayr M: Prevalence and impact of abnormal ROTEM(R) assays in severe blunt trauma: results of the 'Diagnosis and Treatment of Trauma-Induced Coagulopathy (DIA-TRE-TIC) study'. *Br J Anaesth* 2011;107:378–387.
- 45 Casutt M, Kristoff A, Schuepfer G, Spahn DR, Konrad C: Effects on coagulation of balanced (130/0.42) and non-balanced (130/0.4) hydroxyethyl starch or gelatin compared with balanced Ringer's solution: an in vitro study using two different viscoelastic coagulation tests ROTEMTM and SONOCLOTTM. *Br J Anaesth* 2010; 105:273–281.
- 46 Forestier F, Belisle S, Contant C, Harel F, Janvier G, Hardy JF: Reproducibility and interchangeability of the Thromboelastograph, Sonoclot and Hemochron activated coagulation time in cardiac surgery (in French). *Can J Anaesth* 2001;48:902–910.
- 47 Frith D, Goslings JC, Gaarder C, Maegele M, Cohen MJ, Allard S, Johansson PI, Stanworth S, Thiemermann C, Brohi K: Definition and drivers of acute traumatic coagulopathy: clinical and experimental investigations. *J Thromb Haemost* 2010;8:1919–1925.
- 48 Davenport RA, Guerreiro M, Frith D, Rourke C, Platten S, Cohen M, Pearce R, Thiemermann C, Brohi K: Activated protein C drives the hyperfibrinolysis of acute traumatic coagulopathy. *Anesthesiology* 2017; 126:115–127.
- 49 Peltan ID, Rowhani-Rahbar A, Vande Vusse LK, Caldwell E, Rea TD, Maier RV, Watkins TR: Development and validation of a prehospital prediction model for acute traumatic coagulopathy. *Crit Care* 2016;20:371.